

On the Chung-Diaconis-Graham random process

Martin Hildebrand
 Department of Mathematics and Statistics
 University at Albany
 State University of New York
 Albany, NY 12222

February 2, 2008

Abstract

Chung, Diaconis, and Graham considered random processes of the form $X_{n+1} = 2X_n + b_n \pmod{p}$ where $X_0 = 0$, p is odd, and b_n for $n = 0, 1, 2, \dots$ are i.i.d. random variables on $\{-1, 0, 1\}$. If $\Pr(b_n = -1) = \Pr(b_n = 1) = \beta$ and $\Pr(b_n = 0) = 1 - 2\beta$, they asked which value of β makes X_n get close to uniformly distributed on the integers mod p the slowest. In this paper, we extend the results of Chung, Diaconis, and Graham in the case $p = 2^t - 1$ to show that for $0 < \beta \leq 1/2$, there is no such value of β .

1 Introduction

In [1], Chung, Diaconis, and Graham considered random processes of the form $X_{n+1} = 2X_n + b_n \pmod{p}$ where p is an odd integer, $X_0 = 0$, and $b_0, b_1, b_2, \dots$ are i.i.d. random variables. This process is also described in Diaconis [2], and generalizations involving random processes of the form $X_{n+1} = a_n X_n + b_n \pmod{p}$ where (a_i, b_i) for $i = 0, 1, 2, \dots$ are i.i.d. were considered by the author in [3] and [4]. A question asked in [1] concerns cases where $\Pr(b_n = 1) = \Pr(b_n = -1) = \beta$ and $\Pr(b_n = 0) = 1 - 2\beta$. If $\beta = 1/4$ or $\beta = 1/2$, then P_n is close to the uniform distribution (in variation distance) on the integers mod p if n is a large enough multiple of $\log p$ where $P_n(s) = \Pr(X_n = s)$. If $\beta = 1/3$, however, for n a small enough multiple of $(\log p) \log(\log p)$, the variation distance $\|P_n - U\|$ is far from 0 for certain values of p such as $p = 2^t - 1$. Chung, Diaconis, and Graham comment “It would be interesting to know which value of β maximizes the value of N required for $\|P_N - U\| \rightarrow 0$.”

If $\beta = 0$, then $X_n = 0$ with probability 1 for all n . Thus we shall only consider the case $\beta > 0$. We shall show that unless $\beta = 1/4$ or $\beta = 1/2$, then there exists a value $c_\beta > 0$ such that for certain values of p (namely $p = 2^t - 1$), if $n \leq c_\beta (\log p) \log(\log p)$, then $\|P_n - U\| \rightarrow 1$ as $t \rightarrow \infty$. Furthermore, one can have $c_\beta \rightarrow \infty$ as $\beta \rightarrow 0^+$. Work of the author [3] shows that for each β , there is

a value c'_β such that if $n \geq c'_\beta(\log p) \log(\log p)$, then $\|P_n - U\| \rightarrow 0$ as $p \rightarrow \infty$. Thus one may conclude that there is no value of β which maximizes the value of N required for $\|P_N - U\| \rightarrow 0$.

This paper will consider a broader class of distributions for b_n . In particular, $\Pr(b_n = 1)$ need not equal $\Pr(b_n = -1)$. The main argument here relies on a generalization of an argument in [1].

2 Notation and Main Theorem

Recall that the variation distance of a probability P on a finite group G from the uniform distribution on G is given by

$$\begin{aligned} \|P - U\| &= \frac{1}{2} \sum_{s \in G} |P(s) - 1/|G|| \\ &= \max_{A \subseteq G} |P(A) - U(A)| \\ &= \sum_{s: P(s) > 1/|G|} |P(s) - 1/|G|| \end{aligned}$$

The following assumptions are used in the main theorem. Suppose $\Pr(b_n = 1) = a$, $\Pr(b_n = 0) = b$, and $\Pr(b_n = -1) = c$. We assume $a + b + c = 1$ and a , b , and c are all less than 1. Suppose $b_0, b_1, b_2, \dots$ are i.i.d. and $X_0 = 0$. Suppose $X_{n+1} = 2X_n + b_n \pmod{p}$ and p is odd. Let $P_n(s) = \Pr(X_n = s)$. The theorem itself follows:

Theorem 1 *Case 1: Suppose either $b = 0$ and $a = c = 1/2$ or $b = 1/2$. If $n > c_1 \log_2 p$ where $c_1 > 1$ is constant, then $\|P_n - U\| \rightarrow 0$ as $p \rightarrow \infty$ where p is an odd integer.*

Case 2: Suppose a , b , and c do not satisfy the conditions in Case 1. Then there exists a value c_2 (depending on a , b , and c) such that if $n < c_2(\log p) \log(\log p)$ and $p = 2^t - 1$, then $\|P_n - U\| \rightarrow 1$ as $t \rightarrow \infty$.

3 Proof of Case 1

First let's consider the case where $b = 1/2$. Then $b_n = e_n + d_n$ where e_n and d_n are independent random variables with $\Pr(e_n = 0) = \Pr(e_n = 1) = 1/2$, $\Pr(d_n = -1) = 2c$, and $\Pr(d_n = 0) = 2a$. (Note that here $a + c = 1/2 = b$. Thus $2a + 2c = 1$.) Observe that

$$\begin{aligned} X_n &= \sum_{j=0}^{n-1} 2^{n-1-j} b_j \pmod{p} \\ &= \sum_{j=0}^{n-1} 2^{n-1-j} e_j + \sum_{j=0}^{n-1} 2^{n-1-j} d_j \pmod{p} \end{aligned}$$

Let

$$Y_n = \sum_{j=0}^{n-1} 2^{n-1-j} e_j \pmod{p}.$$

If P_n is the probability distribution of X_n (i.e. $P_n(s) = \Pr(X_n = s)$) and Q_n is the probability distribution of Y_n , then the independence of e_n and d_n implies $\|P_n - U\| \leq \|Q_n - U\|$. Observe that on the integers, $\sum_{j=0}^{n-1} 2^{n-1-j} e_j$ is uniformly distributed on the set $\{0, 1, \dots, 2^n - 1\}$. Each element of the integers mod p appears either $\lfloor 2^n/p \rfloor$ times or $\lceil 2^n/p \rceil$ times. Thus

$$\|Q_n - U\| \leq p \left(\frac{\lceil 2^n/p \rceil}{2^n} - \frac{1}{p} \right) \leq \frac{p}{2^n}.$$

If $n > c_1 \log_2 p$ where $c_1 > 1$, then $2^n > p^{c_1}$ and $\|Q_n - U\| \leq 1/p^{c_1-1} \rightarrow 0$ as $p \rightarrow \infty$.

The case where $b = 0$ and $a = c = 1/2$ is alluded to in [1] and left as an exercise. $\square$

4 Proof of Case 2

The proof of this case follows the proof of Theorem 2 in [1] with some modifications. First we assume $b > 0$.

Define, as in [1], the separating function $f : \mathbb{Z}/p\mathbb{Z} \rightarrow \mathbb{C}$ by

$$f(k) := \sum_{j=0}^{t-1} q^{k2^j}$$

where $q := q(p) := e^{2\pi i/p}$. We shall suppose $n = rt$ where r is an integer of the form $r = \delta \log t - d$ for a fixed value δ .

If $0 \leq j \leq t-1$, define

$$\Pi_j := \prod_{\alpha=0}^{t-1} \left(a q^{(2^\alpha(2^j-1))} + b + c q^{-(2^\alpha(2^j-1))} \right).$$

Note that if $a = b = c = 1/3$, then this expression is the same as Π_j defined in the proof of Theorem 2 in [1].

As in the proof of Theorem 2 in [1], $E_U(f) = 0$ and $E_U(f\bar{f}) = t$. Furthermore

$$\begin{aligned} E_{P_n}(f) &= \sum_k P_n(k) f(k) \\ &= \sum_k \sum_{j=0}^{t-1} P_n(k) q^{k2^j} \\ &= \sum_{j=0}^{t-1} \hat{P}_n(2^j) \end{aligned}$$

$$\begin{aligned}
&= \sum_{j=0}^{t-1} \prod_{\alpha=0}^{t-1} \left(aq^{2^\alpha 2^j/p} + b + cq^{-2^\alpha 2^j/p} \right)^r \\
&= t\Pi_1^r.
\end{aligned}$$

Also note

$$\begin{aligned}
E_{P_n}(f\bar{f}) &= \sum_k P_n(k) f(k) \bar{f}(k) \\
&= \sum_k \sum_{j,j'} P_n(k) q^{k(2^j - 2^{j'})} \\
&= \sum_{j,j'} \hat{P}_n(2^j - 2^{j'}) \\
&= \sum_{j,j'} \prod_{\alpha=0}^{t-1} \left(aq^{2^\alpha(2^j - 2^{j'})} + b + cq^{-2^\alpha(2^j - 2^{j'})} \right)^r \\
&= t \sum_{j=0}^{t-1} \Pi_j^r.
\end{aligned}$$

(Note that the expressions for $E_{P_N}(f)$ and $E_{P_N}(f\bar{f})$ in the proof of Theorem 2 of [1] have some minor misprints.)

The (complex) variances of f under U and P_n are $\text{Var}_U(f) = t$ and

$$\begin{aligned}
\text{Var}_{P_n}(f) &= E_{P_n}(|f - E_{P_n}(f)|^2) \\
&= E_{P_n}(f\bar{f}) - E_{P_n}(f)E_{P_n}(\bar{f}) \\
&= t \sum_{j=0}^{t-1} \Pi_j^r - t^2 |\Pi_1|^{2r}.
\end{aligned}$$

Like [1], we use the following complex form of Chebyshev's inequality for any Q :

$$Q \left(\left\{ x : |f(x) - E_Q(f)| \geq \alpha \sqrt{\text{Var}_Q(f)} \right\} \right) \leq 1/\alpha^2$$

where $\alpha > 0$. Thus

$$U \left(\left\{ x : |f(x)| \geq \alpha t^{1/2} \right\} \right) \leq 1/\alpha^2$$

and

$$P_n \left(\left\{ x : |f(x) - t\Pi_1^r| \geq \beta \left(t \sum_{j=0}^{t-1} \Pi_j^r - t^2 |\Pi_1|^{2r} \right)^{1/2} \right\} \right) \leq 1/\beta^2.$$

Let A and B denote the complements of these 2 sets; thus $U(A) \geq 1 - 1/\alpha^2$ and $P_n(B) \geq 1 - 1/\beta^2$. If A and B are disjoint, then $\|P_n - U\| \geq 1 - 1/\alpha^2 - 1/\beta^2$.

Suppose r is an integer with

$$r = \frac{\log t}{2 \log(1/|\Pi_1|)} - \lambda$$

where $\lambda \rightarrow \infty$ as $t \rightarrow \infty$ but $\lambda \ll \log t$. Then $t|\Pi_1|^r = t^{1/2}|\Pi_1|^{-\lambda} \gg t^{1/2}$. Observe that the fact a , b , and c do not satisfy the conditions in Case 1 implies $|\Pi_1|$ is bounded away from 0 as $t \rightarrow \infty$. Furthermore $|\Pi_1|$ is bounded away from 1 for a given a , b , and c .

In contrast, let's consider what happens to $|\Pi_1|$ if a , b , and c do satisfy the condition in Case 1. If $b = 1/2$, then the $\alpha = t - 1$ term in the definition of Π_1 converges to 0 as $t \rightarrow \infty$ and thus Π_1 also converges to 0 as $t \rightarrow \infty$ since each other term has length at most 1. If $a = c = 1/2$ and $b = 0$, then the $\alpha = t - 2$ term in the definition of Π_1 converges to 0 as $t \rightarrow \infty$ and thus Π_1 also converges to 0 as $t \rightarrow \infty$.

Claim 1

$$\frac{1}{t} \sum_{j=0}^{t-1} \left(\frac{\Pi_j}{|\Pi_1|^2} \right)^r \rightarrow 1$$

as $t \rightarrow \infty$.

Note that this claim implies $(\text{Var}_{P_n}(f))^{1/2} = o(E_{P_n}(f))$ and thus Case 2 of Theorem 1 follows.

Note that $\Pi_0 = 1$. By Proposition 1 below, $\bar{\Pi}_j = \Pi_{t-j}$. Thus $t \sum_{j=0}^{t-1} \Pi_j^r$ is real. Also note that since $\text{Var}_{P_n}(f) \geq 0$, we have

$$\frac{t \sum_{j=0}^{t-1} \Pi_j^r}{t^2 |\Pi_1|^{2r}} \geq 1.$$

Thus to prove the claim, it suffices to show

$$\frac{1}{t} \sum_{j=0}^{t-1} \left(\frac{|\Pi_j|}{|\Pi_1|^2} \right)^r \rightarrow 1.$$

Proposition 1 $\bar{\Pi}_j = \Pi_{t-j}$.

Proof: Note that

$$\bar{\Pi}_j = \prod_{\alpha=0}^{t-1} \left(aq^{-(2^\alpha(2^j-1))} + b + cq^{(2^\alpha(2^j-1))} \right)$$

and

$$\Pi_{t-j} = \prod_{\beta=0}^{t-1} \left(aq^{(2^\beta(2^{t-j}-1))} + b + cq^{-(2^\beta(2^{t-j}-1))} \right).$$

If $j \leq \beta \leq t-1$, then note

$$\begin{aligned} 2^\beta(2^{t-j}-1) &= 2^{\beta-j}(2^t-2^j) \\ &= 2^{\beta-j}(1-2^j) \pmod{p} \\ &= -2^{\beta-j}(2^j-1). \end{aligned}$$

Thus the terms in Π_{t-j} with $j \leq \beta \leq t-1$ are equal to the terms in $\overline{\Pi}_j$ with $0 \leq \alpha \leq t-j-1$.

If $0 \leq \beta \leq j-1$, then note

$$\begin{aligned} 2^\beta(2^{t-j}-1) &= 2^{t+\beta}(2^{t-j}-1) \pmod{p} \\ &= 2^{t+\beta-j}(2^t-2^j) \\ &= 2^{t+\beta-j}(1-2^j) \pmod{p} \\ &= -2^{t+\beta-j}(2^j-1). \end{aligned}$$

Thus the terms in Π_{t-j} with $0 \leq \beta \leq j-1$ are equal to the terms in $\overline{\Pi}_j$ with $t-j \leq \alpha \leq t-1$. $\square$

Now let's prove the claim. Let $G(x) = |ae^{2\pi ix} + b + ce^{-2\pi ix}|$. Thus

$$|\Pi_j| = \prod_{\alpha=0}^{t-1} G(2^\alpha(2^j-1)/p).$$

Note that if $0 \leq x < y \leq 1/4$, then $G(x) > G(y)$. On the interval $[1/4, 1/2]$, where G increases and where G decreases depends on a , b , and c .

We shall prove a couple of facts analogous to facts in [1].

Fact 1: There exists a value t_0 (possibly depending on a , b , and c) such that if $t > t_0$, then $|\Pi_j| \leq |\Pi_1|$ for all $j \geq 1$.

Since $G(x) = G(1-x)$, in proving this fact we may assume without loss of generality that $2 \leq j \leq t/2$. Note that

$$|\Pi_j| = \prod_{i=0}^{t-j-1} G\left(\frac{2^{i+j}-2^i}{p}\right) \prod_{i=0}^{j-1} G\left(\frac{2^{i+t-j}-2^i}{p}\right).$$

We associate factors x from $|\Pi_j|$ with corresponding factors $\pi(x)$ of $|\Pi_1|$ in a manner similar to that in [1]. For $0 \leq i \leq t-j-2$, associate $G((2^{i+j}-2^i)/p)$ with $G(2^{i+j-1}/p)$. Note that for $0 \leq i \leq t-j-2$, we have $G((2^{i+j}-2^i)/p) \leq G(2^{i+j-1}/p)$. For $0 \leq i \leq j-3$, associate $G((2^{i+t-j}-2^i)/p)$ in $|\Pi_j|$ with $G(2^i/p)$ in $|\Pi_1|$. Note that for $0 \leq i \leq j-3$, we have $G((2^{i+t-j}-2^i)/p) \leq G(2^i/p)$.

The remaining terms in $|\Pi_j|$ are

$$G\left(\frac{2^{t-1}-2^{t-j-1}}{p}\right) G\left(\frac{2^{t-1}-2^{j-1}}{p}\right) G\left(\frac{2^{t-2}-2^{j-2}}{p}\right)$$

and the remaining terms in $|\Pi_1|$ are

$$G\left(\frac{2^{t-1}}{p}\right) G\left(\frac{2^{t-2}}{p}\right) G\left(\frac{2^{j-2}}{p}\right).$$

It can be shown that since $b > 0$,

$$\limsup_{t \rightarrow \infty} \frac{G\left(\frac{2^{t-1}-2^{t-j-1}}{p}\right) G\left(\frac{2^{t-1}-2^{j-1}}{p}\right) G\left(\frac{2^{t-2}-2^{j-2}}{p}\right)}{G\left(\frac{2^{t-1}}{p}\right) G\left(\frac{2^{t-2}}{p}\right) G\left(\frac{2^{j-2}}{p}\right)} \leq \frac{M}{G(0)} < 1$$

where $M = \sup_{x \in [1/4, 1/2]} G(x)$. Note $G(0) = 1$. To see the fact that $M < 1$ if $b > 0$, consider $H(x) = |ae^{2\pi i x} + b|$ if $a > 0$. $H(x)$ is decreasing on $[0, 1/2]$, and $a + b - H(1/4) > 0$. For $x \in [1/4, 1/2]$, we have $G(x) \leq H(x) + c = 1 - (a + b - H(x)) \leq 1 - (a + b - H(1/4))$. If $a = 0$, then $c > 0$ and a similar argument involving $|ce^{-2\pi i x} + b|$ applies. Indeed, for some t_0 , if $t > t_0$ and $2 \leq j \leq t/2$,

$$\begin{aligned} & G\left(\frac{2^{t-1}-2^{t-j-1}}{p}\right) G\left(\frac{2^{t-1}-2^{j-1}}{p}\right) G\left(\frac{2^{t-2}-2^{j-2}}{p}\right) \\ & \leq G\left(\frac{2^{t-1}}{p}\right) G\left(\frac{2^{t-2}}{p}\right) G\left(\frac{2^{j-2}}{p}\right). \end{aligned}$$

□

Fact 2: There exists a value t_1 (possibly depending on a , b , and c) such that if $t > t_1$, then the following holds. There is a constant c_0 such that for $t^{1/3} \leq j \leq t/2$, we have

$$\frac{|\Pi_j|}{|\Pi_1|^2} \leq 1 + \frac{c_0}{2^j}$$

To prove this fact, we associate, for $i = 0, 1, \dots, j-1$, the terms

$$G\left(\frac{2^{t-i-1}-2^{j-i-1}}{p}\right) G\left(\frac{2^{t-i-1}-2^{t-j-i-1}}{p}\right)$$

in $|\Pi_j|$ with the terms

$$\left(G\left(\frac{2^{t-i-1}}{p}\right)\right)^2$$

in $|\Pi_1|^2$. Suppose $A = \max |G'(x)|$. Note that $A < \infty$. Then

$$\left|G\left(\frac{2^{t-i-1}-2^{j-i-1}}{p}\right)\right| \leq \left|G\left(\frac{2^{t-i-1}}{p}\right)\right| + A \frac{2^{j-i-1}}{p}.$$

Thus

$$\frac{\left|G\left(\frac{2^{t-i-1}-2^{j-i-1}}{p}\right)\right|}{\left|G\left(\frac{2^{t-i-1}}{p}\right)\right|} \leq 1 + A \frac{2^{j-i-1}}{p \left|G\left(\frac{2^{t-i-1}}{p}\right)\right|}.$$

Likewise

$$\frac{\left|G\left(\frac{2^{t-i-1}-2^{t-j-i-1}}{p}\right)\right|}{\left|G\left(\frac{2^{t-i-1}}{p}\right)\right|} \leq 1 + A \frac{2^{t-j-i-1}}{p \left|G\left(\frac{2^{t-i-1}}{p}\right)\right|}.$$

Since we do not have the conditions for Case 1, there is a positive value B and value t_2 such that if $t > t_2$, then $|G(2^{t-i-1}/p)| > B$ for all i with $0 \leq i \leq j-1$. By an exercise, one can verify

$$\prod_{i=0}^{j-1} \frac{\left| G\left(\frac{2^{t-i-1}-2^{j-i-1}}{p}\right) G\left(\frac{2^{t-i-1}-2^{t-j-i-1}}{p}\right) \right|}{\left| G\left(\frac{2^{t-i-1}}{p}\right) \right|^2} \leq 1 + \frac{c_3}{2^j}$$

for some value c_3 not depending on j .

Note that the remaining terms in $|\Pi_j|$ all have length less than 1. The remaining terms in $|\Pi_1|^2$ are

$$\prod_{i=j}^{t-1} \left| G\left(\frac{2^{t-i-1}}{p}\right) \right|^2.$$

Since $G'(0) = 0$, there are positive constants c_4 and c_5 such that

$$\left| G\left(\frac{2^{t-i-1}}{p}\right) \right| \geq 1 - c_4 \left(\frac{2^{t-i-1}}{p}\right)^2 \geq \exp\left(-c_5 \frac{2^{t-i-1}}{p}\right)$$

for $i \geq j \geq t^{1/3}$. Observe

$$\begin{aligned} \prod_{i=j}^{t-1} \exp\left(-c_5 \frac{2^{t-i-1}}{p}\right) &= \exp\left(-c_5 \sum_{i=j}^{t-1} 2^{t-i-1}/p\right) \\ &= \exp\left(-c_5 \sum_{k=0}^{t-j-1} 2^k/p\right) \\ &= \exp\left(-c_5 \frac{2^{t-j} - 1}{2^t - 1}\right) \\ &> \exp\left(-c_5 \frac{2^{t-j}}{2^t}\right) \\ &= \exp(-c_5/2^j) > 1 - c_5/2^j. \end{aligned}$$

There exists a constant c_0 such that

$$\frac{1 + c_3/2^j}{(1 - c_5/2^j)^2} \leq 1 + c_0/2^j$$

for $j \geq 1$.

Thus, as in [1],

$$\sum_{t^{1/3} \leq j \leq t/2} \left| \left(\frac{|\Pi_j|}{|\Pi_1|^2} \right)^r - 1 \right| \leq \frac{c_6 t r}{2^{t^{1/3}}} < \frac{c_7}{2^{t^{1/4}}}$$

for values c_6 and c_7 . Since $|\Pi_j| = |\Pi_{t-j}|$,

$$\begin{aligned} \frac{1}{t} \sum_{j=0}^{t-1} \left(\frac{|\Pi_j|}{|\Pi_1|^2} \right)^r &\leq \frac{1}{t} \frac{1}{|\Pi_1|^{2r}} + \frac{2}{t} \left(\sum_{1 \leq j < t^{1/3}} \left(\frac{|\Pi_j|}{|\Pi_1|^2} \right)^r + \sum_{t^{1/3} \leq j \leq t/2} \left(\frac{|\Pi_j|}{|\Pi_1|^2} \right)^r \right) \\ &= 1 + o(1) \end{aligned}$$

as $t \rightarrow \infty$. Thus Fact 2, the claim, and Theorem 1 if $b > 0$ are proved.

If $b = 0$, Theorem 1 can be proved by considering the following random processes on the integers mod p :

1. $X_0 = 0$ and $X_{n+1} = 2X_n + b_n \pmod{p}$ where $P(b_n = 1) = a$ and $P(b_n = -1) = 1 - a$
2. $Y_0 = 0$ and $Y_{n+1} = 2Y_n + d_n \pmod{p}$ where $P(d_n = 2) = a$ and $P(d_n = 0) = 1 - a$
3. $Z_0 = 0$ and $Z_{n+1} = 2Z_n + e_n \pmod{p}$ where $P(e_n = 1) = a$ and $P(e_n = 0) = 1 - a$

If $P_n(s) = \Pr(X_n = s)$, $Q_n(s) = \Pr(Y_n = s)$, and $R_n(s) = \Pr(Z_n = s)$, then $\|P_n - U\| = \|Q_n - U\| = \|R_n - U\|$. To see this, we can let $d_n = 2e_n$ and $b_n = d_n - 1$ so that $Y_n = 2Z_n$ and $X_n = Y_n - \sum_{j=0}^{n-1} 2^j$ for $n \geq 1$. To conclude the case where $b = 0$, use the argument with $b \neq 0$ on the third random process (provided that $a \neq 1/2$ so that we are in Case 2). $\square$

The next proposition considers what happens as we vary the values a , b , and c .

Proposition 2 *If $a = c = \beta$ and $b = 1 - 2\beta$ and $m_\beta = \liminf_{t \rightarrow \infty} |\Pi_1|$, then $\lim_{\beta \rightarrow 0^+} m_\beta = 1$.*

Proof: Suppose $\beta < 1/4$. Then

$$\Pi_1 = \prod_{\alpha=0}^{t-1} ((1 - 2\beta) + 2\beta \cos(2\pi 2^\alpha / p)).$$

Let $h(\alpha) = (1 - 2\beta) + 2\beta \cos(2\pi 2^\alpha / p)$. Note that

$$\begin{aligned} \lim_{\beta \rightarrow 0^+} h(t-1) &= 1 \\ \lim_{\beta \rightarrow 0^+} h(t-2) &= 1 \\ \lim_{\beta \rightarrow 0^+} h(t-3) &= 1 \end{aligned}$$

Furthermore, for some constant $\gamma > 0$, one can show

$$h(\alpha) > \exp(-\beta\gamma(2^\alpha/p)^2)$$

if $2^\alpha/p \leq 1/8$ and $0 < \beta < 1/10$. So

$$\prod_{\alpha=0}^{t-4} h(\alpha) > \prod_{\alpha=0}^{t-4} \exp(-\beta\gamma(2^\alpha/p)^2)$$

$$\begin{aligned}
&= \exp\left(-\beta\gamma\sum_{\alpha=0}^{t-4}(2^\alpha/p)^2\right) \\
&> \exp(-\beta\gamma 2^{2(t-4)}(4/3)/p^2) \rightarrow 1
\end{aligned}$$

as $\beta \rightarrow 0^+$.

□

Recalling that

$$r = \frac{\log t}{2 \log(1/|\Pi_1|)} - \lambda,$$

we see that $1/(2 \log(1/|\Pi_1|))$ can be made arbitrarily large by choosing β small enough. Thus there exist values $c_\beta \rightarrow \infty$ as $\beta \rightarrow 0^+$ such that if $n \leq c_\beta(\log p) \log(\log p)$, then $\|P_n - U\| \rightarrow 1$ as $t \rightarrow \infty$.

5 Problems for further study

One possible problem is to see if in some sense, there is a value of β on $[1/4, 1/2]$ which maximizes the value of N required for $\|P_N - U\| \rightarrow 0$; to consider such a question, one might restrict p to values such that $p = 2^t - 1$.

Another possible question considers the behavior of these random processes for almost all odd p . For $\beta = 1/3$, Chung, Diaconis, and Graham [1] showed that a multiple of $\log p$ steps suffice for almost all odd p . While their arguments should be adaptable with the change of appropriate constants to a broad range of choices of a , b , and c in Case 2, a more challenging question is to determine for which a , b , and c in Case 2 (if any), $(1 + o(1)) \log_2 p$ steps suffice for almost all odd p .

6 Acknowledgments

The author thanks Ron Graham for mentioning the problem at a January, 2005, conference on the Mathematics of Persi Diaconis. The author also thanks Robin Pemantle for a conversation on the topic and the participants in the Probability and Related Fields Seminar at the University at Albany for listening to some ideas on the problem. The author also thanks Ravi Montenegro for pointing out a flaw in the original argument, which appeared in [5].

References

- [1] Chung, F., Diaconis, P., and Graham, R. A random walk problem arising in random number generation. *Ann. Probab.* **15** (1987), 1148-1165.
- [2] Diaconis, P. *Group Representations in Probability and Statistics*. Institute of Mathematical Statistics, 1988.
- [3] Hildebrand, M. Random processes of the form $X_{n+1} = a_n X_n + b_n \pmod{p}$. *Ann. Probab.* **21** (1993), 710-720.

- [4] Hildebrand, M. Random processes of the form $X_{n+1} = a_n X_n + b_n \pmod{p}$ where b_n takes on a single value. pp. 153-174, *Random Discrete Structures*, ed. Aldous and Pemantle. Springer-Verlag, 1996.
- [5] Hildebrand, M. On the Chung-Diaconis-Graham random process. *Electron. Comm. in Probab.* **11** (2006), 347-356.